

Collembola Diversity of Moss Habitats in the Sopron Region, NW-Hungary

György TRASER^{a*} – Péter SZÜCS^b – Dániel WINKLER^c

^a Institute of Forest and Wood Protection, University of West Hungary, Sopron, Hungary

^b Institute of Botany, University of West Hungary, Sopron, Hungary

^c Institute of Wildlife Management, University of West Hungary, Sopron, Hungary

Abstract – The *Collembola* fauna of the moss flora in the Sopron region was studied. 18 moss species as well as 3.451 *Collembola* specimens (belonging to 60 species) were collected in 2003/04 in moss samples of three habitats. The highest *Collembola* diversity was found in Tómalom (a reed bed habitat) where very low abundance and relatively high species richness were characteristic. The *Collembola* diversity of the other two habitats (Sopron, Botanic Garden; Fertőrákos – a dry xerophile grass habitat) was lower. The results have shown a relatively high similarity between the *Collembola* communities in Sopron and Fertőrákos while just the opposite was observed between the Sopron and Tómalom samples.

bryofauna / Collembola diversity / dominance structure

Kivonat – Sopron környéki mohás élőhelyek *Collembola*-diverzitása. A Sopron környéki mohaflóra feltárása során vizsgáltuk a mohapárnákban élő ugróvillás közösségek diverzitását. 2003/2004-ben három vizsgálati területen gyűjtve 18 mohafaj mintáiban összesen 60 faj 3451 egyede került elő. A legmagasabb diverzitást a Tómalomnál, egy nádas-vizes habitatban találtuk. A két másik területen (Sopron, Botanikus kert; Fertőrákos, száraz sziklagyep) az egyedszám magasabb volt, a diverzitás azonban alacsonyabbnak adódott. Az eredmények a *Collembola* közösségek viszonylag nagy hasonlóságát mutatták Sopron és Fertőrákos viszonylatában, míg ennek ellenkezője igaz Sopron és Tómalom között.

bryofauna / Collembola-diverzitás / dominancia-viszonyok

1 INTRODUCTION

The *Collembola* (springtails) are the most abundant *Apterygota Entognatha* insects in the world, found in vast numbers from the tropics to the poles. The *Collembola* are mostly connected to forest habitats. They are chiefly soil dwellers, but we can also find springtails living in the canopy of tropical rain forests and they are also characteristic members of the bryofauna (Varga – Vargha 1992, Varga – Oldal 1998, Palacios-Vargas et al. 1998, 1999). In individual cases *Collembola* density can reach 98% proportion of the total density of *Arthropoda* collected. An average square metre of soil in a temperate woodland can yield up to 50.000 individuals (Dunger 1983). Moreover, Dunger (1983) reveals that the optimal *Collembola* density would be tenfold of the mentioned value.

* Corresponding author: traser@emk.nyme.hu; H-9400 SOPRON, Bajcsy-Zs. u. 4.

The ecological importance of Collembola is indispensable in the decomposition of litter. They have an important role also in the food chain since they often form an important food source for predator Arthropods (e.g. for ground beetles). Most Collembola species prefer humid habitats. Their occurrence is furthermore determined by vegetation types, so springtails are important ecological indicator species for habitats and microhabitats. Although a paper on this region's Collembola fauna has already been published (Traser 2002), it doesn't treat the aspect of the springtail-moss connection.

The aim of this study was to determine the relationship between Collembola community composition, species richness, density, diversity in three different moss habitats in the Sopron region. Further investigations were related to the comparison of structural properties of Collembola communities between the moss habitats.

2 MATERIAL AND METHODS

2.1 Study area

The heart of the Sopron region is formed by the valley of the Ikva streamlet between the Sopron Mountains and the Szárhalmi Hillocks. The three study areas are the following: the Botanic Garden (Sopron), a wooded area with grass mosaic; Tómalom, a reed bed habitat and Fertőrákos, a dry limestone/grass habitat. *Table 1* shows further details on the study sites.

Table 1. Moss samples data and locality

Location	Moss samples in 2003 / 2004		
	Date	GPS coordinate	Habitat
Sopron – Botanic Garden	06. 03. 2003.	47.68056°N	wooded area with grass mosaic
	09. 10. 2003.	16.58222°E	
	02. 03. 2004.		
Tómalom – reed bed	10. 09. 2003.	47.70965°N	<i>Caricion davallianae</i> reed, wet area
		16.62362°E	
Fertőrákos – dry grass	06. 05. 2004.	47.72844°N	<i>Festucetalia valesiacae</i> limestone grassland
		16.64010°E	

2.2 Data collection and analysis

Samples of 100 cm³ of moss were collected. A total of 43 samples were obtained from the three different habitats ranging from 10 to 19 samples per site (Tómalom:10; Sopron: 14; Fertőrákos: 19).

Extraction of *Collembola* specimens from the moss was carried with the help of a modified Tullgren's apparatus at room temperature (Balogh 1958). The *Collembola* and moss species have been stored in the archives of the authors. For identification of the *Collembola* fauna we used the keys of Gisin (1960), Bretfeld (1999), Jordana et al. (1997), Babenko et al.(1994), Deharveng (1982), Fjellberg (1980, 1998), Massoud (1967), Potapow (2001), Pomorski (1998), Thibaud et al. (2004), Zimdars & Dunger (1994), Stach (1960, 1963), Weiner (1996) and several other papers, while for the moss flora we used the keys of Orbán – Vajda (1983), Smith (1978, 1990) and Hedenas (1993).

Graphical methods (dominance curves) were employed to allow direct comparison of *Collembola* species diversity characteristics between the sites.

To measure *Collembola* similarity (based on dominance) between the habitats the Renkonen coefficient (Re %) was used:

$$\text{Re \%} = \sum \min D_{A:B}$$

where “D_A“ and “D_B”: dominance % of the common species in habitat “A” and “B”

Another similarity coefficients such as the Sørensen quotient ($QS\% = 2G / (S_A + S_B) * 100$), the Jaccard’s species identity index [$J_N\% = G * 100 / (S_A + S_B - G)$] and the similarity index of Weinstein ($K_w = Re * J_N$) were calculated to determine the similarities/differences between the habitats and their collembolan communities. (Symbols are: “G”= number of the common species in the paired habitats; “S_A“ and “S_B“ are the species numbers in habitat “A” and habitat “B”).

To compare *Collembola* diversity values of two different habitats a *t* test was used.

3 RESULTS AND DISCUSSION

Table 2 shows the distribution and quantity of the moss samples in the three different habitats. There are no common moss species in these habitats except for two species that are present in both Sopron and Tómalom and one species found both in Sopron and Fertőrákos.

Table 2. Moss species and the number of samples taken

Moss species	Sopron	Tómalom	Fertőrákos
<i>Tortella tortuosa</i> (HEDW.) LIMPR.			3
<i>Pleurochaete squarrosa</i> (BRID.) LINDB.			4
<i>Plagiomnium elatum</i> (BRUCH et SCHIMP.) T. J. KOP.		1	
<i>Plagiomnium undulatum</i> (HEDW.) T. J. KOP.	2		
<i>Climacium dendroides</i> (HEDW.) F. WEBER et D. MOHR	1		
<i>Thuidium philibertii</i> LIMPR.	1		
<i>Thuidium abietinum</i> (HEDW.) SCHIMP.			4
<i>Campylium chrysophyllum</i> (BRID.) LANGE			1
<i>Amblystegium varium</i> (HEDW.) LINDB.		2	
<i>Drepanocladus cossonii</i> (SCHIMP.) LOESKE		2	
<i>Calliergonella cuspidata</i> (HEDW.) LOESKE	3	4	
<i>Homalothecium lutescens</i> (HEDW.) H. ROB			3
<i>Brachythecium rutabulum</i> (HEDW.) SCHIMP.	1	1	
<i>Scleropodium purum</i> (HEDW.) LIMPR.	2		
<i>Eurhynchium hians</i> (HEDW.) SANDLE LAC.	1		
<i>Hypnum cupressiforme</i> HEDW.	2		3
<i>Rhytidium rugosum</i> (HEDW.) KINDB.			1
<i>Rhytidiadelphus triquetrus</i> (HEDW.) WARNST.	1		
number of moss samples	14	10	19
moss species number	9	5	7

The moss species *Calliergonella cuspidata* and *Brachythecium rutabulum* were present both in Sopron and Tómalom while *Hypnum cupressiforme* was found both in the Sopron and in the Fertőrákos samples. The common *Collembola* species in the mentioned moss cushions are the followings:

Moss sp.: *Calliergonella cuspidata* (HEDW.) LOESKE

In Tómalom 15 *Collembola* species were collected on September 10, 2003. Out of these species only two spp. (*Sphaeridia pumilis*; *Parisotoma notabilis*) were found common to

both Tómalom and Sopron samples (in Sopron 16 *Collembola* spp. were found on the same moss species).

Moss sp.: *Brachythecium rutabulum* (HEDW.) SCHIMP.

This moss species was collected from monospecific cushions in Tómalom, but mixed with *Scleropodium purum* (HEDW.) LIMPR. and *Eurhynchium hians* (HEDW.) SANDLE LAC. in the Sopron samples. The *Collembola* species richness was 4 in the samples of Tómalom and 9 in Sopron, but no common species to both samples were found.

Moss sp.: *Hypnum cupressiforme* HEDW.

We collected 14 *Collembola* species in the Fertőrákos samples and 9 in the Sopron samples. The only *Collembola* species found common to both samples was *Entomobrya nivalis*.

Table 3 shows the number of *Collembola* individuals in the different habitats. Since the number of samples was not equal in the three habitats, these values can give approximate information on the volume only. The low number of individuals in the Tómalom samples is a very uncommon phenomenon.

Table 3 Number of *Collembola* individuals in the three habitats

	Sopron	Tómalom	Fertőrákos	Σ
Hypogastruridae				
<i>Hypogastrura socialis</i> (Uzel, 1891)	4	0	0	4
<i>Hypogastrura vernalis</i> (Carl, 1901)	0	0	284	284
<i>Xenylla boernerii</i> (Axelson, 1905)	400	0	145	545
<i>Xenylla maritima</i> (Tullberg, 1869)	1	0	0	1
<i>Xenylla brevicauda</i> (Tullberg, 1869)	1	0	120	121
<i>Willemia virae</i> (Kaprus, 1997)	0	0	1	1
Brachystomellidae				
<i>Brachystomella parvula</i> (Schaeffer 1896)	0	10	0	10
Neanuridae				
<i>Friesea truncata</i> (Cassagnau, 1958)	0	10	0	10
<i>Micranurida pygmaea</i> (Börner, 1901)	0	3	0	3
<i>Neanura muscorum</i> (Templeton, 1835)	0	1	0	1
Onychiuridae				
<i>Supraphorura furcifera</i> (Börner, 1901)	0	7	0	7
<i>Protaphorura armata</i> (Tullberg, 1869)	0	0	2	2
<i>Doutnacia xerophila</i> (Rusek, 1974)	4	0	0	4
<i>Mesaphorura critica</i> (Ellis, 1976)	0	0	4	4
<i>Mesaphorura hyliphila</i> (Rusek, 1982)	14	0	0	14
<i>Mesaphorura krausbaueri</i> (Börner, 1901)	0	0	36	36
<i>Mesaphorura macrochaeta</i> (Rusek, 1976)	1	2	0	3
<i>Metaphorura affinis</i> (Börner, 1902)	1	0	0	1
Isotomidae				
<i>Pachyotoma crassicauda</i> (Tullberg, 1871)	0	10	0	10
<i>Cryptopygus bipunctatus</i> (Axelson, 1903)	311	0	319	630
<i>Folsomia manolachei</i> (Bagnall, 1939)	0	0	632	632
<i>Folsomia penicula</i> (Bagnall, 1939)	31	0	0	31
<i>Folsomia quadrioculata</i> (Tullberg, 1871)	0	4	0	4
<i>Isotomiella minor</i> (Schaeffer, 1896)	0	13	2	15
<i>Parisotoma notabilis</i> (Schäffer, 1896)	264	7	159	430
<i>Isotoma viridis</i> (Bourlet, 1839)	0	0	4	4
<i>Isotoma riparia</i> (Nicolet, 1842)	0	1	0	1
<i>Isotomurus cf. palustris</i> (Müller, 1776)	6	0	0	6
<i>Isotomurus prasinus</i> (Reuter, 1891)	0	21	0	21

Table 3 cont. Number of Collembola individuals in the three habitats

	Sopron	Tómalom	Fertőrákos	Σ
Entomobryidae				
<i>Entomobrya corticalis</i> (Nicolet, 1842)	2	0	0	2
<i>Entomobrya handschini</i> (Stach, 1922)	0	0	11	11
<i>Entomobrya multifasciata</i> (Tullberg, 1871)	0	0	4	4
<i>Entomobrya nigriventris</i> (Stach, 1929)	0	0	3	3
<i>Entomobrya nivalis</i> (Linnaeus, 1758)	1	0	11	12
<i>Lepidocyrtus cyaneus</i> (Tullberg, 1871)	0	3	0	3
<i>Lepidocyrtus lanuginosus</i> (Gmelin, 1788)	91	0	25	116
<i>Lepidocyrtus lignorum</i> (Fabricius, 1775)	0	0	2	2
<i>Lepidocyrtus paradoxus</i> (Uzel, 1890)	0	2	0	2
<i>Lepidocyrtus peisonis</i> (Traser et Christian, 1992)	0	15	0	15
<i>Lepidocyrtus violaceus</i> (Fourcroy, 1785)	1	0	0	1
<i>Pseudosinella alba</i> (Packard, 1873)	6	0	1	7
<i>Pseudosinella octopunctata</i> (Börner, 1901)	0	0	32	32
Orchesellidae				
<i>Orchesella cincta</i> (Linnaeus, 1758)	10	0	12	22
<i>Orchesella bifasciata</i> (Nicolet, 1842)	1	0	0	1
<i>Orchesella xerothermica</i> (Stach, 1960)	0	0	2	2
<i>Heteromurus major</i> (Moniez, 1889)	22	0	3	25
<i>Heteromurus nitidus</i> (Templeton, 1835)	11	0	0	11
Tomoceridae				
<i>Tomocerus cf. baudoti</i> (Denis, 1932)	3	0	0	3
<i>Tomocerus minor</i> (Lubbock, 1862)	3	0	0	3
Cyphoderidae				
<i>Cyphoderus albinus</i> (Nicolet, 1842)	1	0	1	2
Oncopoduridae				
<i>Oncopodura crassicornis</i> (Schoebootham, 1911)	0	0	1	1
Neelidae				
<i>Megalothorax minimus</i> (Willem, 1900)	16	13	4	33
<i>Neelides minutus</i> (Folsom, 1901)	0	3	0	3
Sminthuridida				
<i>Sphaeridia pumilis</i> (Krausbauer, 1898)	6	25	0	31
Katiannida				
<i>Sminthurinus elegans</i> (Fitch, 1863)	31	0	0	31
<i>Sminthurinus aureus</i> (Lubbock, 1862)	2	0	0	2
Dicyrtomida				
<i>Dicyrtoma fusca</i> (Lubbock, 1873)	0	2	0	2
Bourletiellida				
<i>Deuterosminthurus bicinctus</i> (Koch, 1840)	0	0	1	1
<i>Fasciosminthurus strigatus</i> (Stach, 1922)	0	1	6	7
<i>Heterosminthurus bilineatus</i> (Bourlet, 1842)	0	0	4	4
Σ	1245	153	1831	3229

Some of the data are interesting since species richness is much higher in the moss cushions of Sopron region than the one recorded in the Bükk Mountains in NE Hungary (Varga 1989, 1991, 1992, 1998).

Collembola dominance curves (Figures 1-3) of the three habitats show very different characteristics. The most abundant species in the samples are different for all three habitats. The dominant species primarily belong to taxa *Isotomidae* and *Hypogastruridae* (followed by taxa *Entomobryidae* and *Symphyleona*). There are only few species belonging to the family

Neanuridae. This proportion of species roughly corresponds to the one characteristic for the Central European *Collembola* fauna.




Figure 1. *Collembola* dominance in Sopron ($n = 1245$ specimens)

The *Collembola* dominance curve of the Sopron sample shows three species with relatively high dominance rank. *Xenylla boernerii*, accounting for > 30 % of the total number of individuals, is a widely distributed species in Europe (including Ukraine and Latvia), occurring in litter, mosses and lichens on trunks and under the bark of trees (Thibaud et al. 2004). *Cryptopygus bipunctatus* ($D = 25\%$) is rather common in Central Europe, rarer in Northern and Southern Europe. It is a xerothermic, ruderal species, mostly occurring in open and disturbed sites (Potapow 2001). *Parisetoma notabilis* ($D = 21\%$) is one of the most ubiquitous *Collembola* of the Western Palaearctic. Its optimum is located in Central Europe and in the European part of Russia. It reaches high densities in both natural and disturbed sites (open grasslands, shelter belts, forests). Its presence in every kind of biotopes (excluding the too extreme ones) is a characteristic of this species (Potapow 2001).




Figure 2. *Collembola* dominance in Tómalom ($n = 153$ specimens)

The most dominant *Collembola* (D = 16,5%) in the Tómalom samples was *Sphaeridia pumilis*, a widespread species in Europe and North America (type locality: Germany, on *Rumex nemorosus* and *Glyceria* of a forest path), occurring on the surface and in upper layers of different moist soil types and in their low vegetation (Bretfeld 1999). Further dominant member (D = 14 %) of this community is *Isotomurus prasinus*, a typical hydrophilous species, common and abundant in various moist sites, often occurring also in agricultural fields (especially in Central Europe). Its exact distribution has still to be determined. Type locality: Siberia, Yenisei River (Potapow 2001). The third most dominant collembola, *Lepidocyrtus peisonis* is a typical hydrophile species, occurring mostly on banks of stagnant eutrophic waters in the Kisalföld region. It has only been recorded in Hungary and Austria (Traser – Christian 1992). Type locality: Fertőrákos, in a humid sedge (*Carex riparia* and *Phragmites australis*) field.




Figure 3. *Collembola* dominance in Fertőrákos (n = 1831 specimens)

The species *Folsomia manolachei* was found the most dominant in the Fertőrákos sample (accounted for 35 % of the total number of individuals). It is a widespread springtail presumably all over the Palaearctic, recorded in almost all European countries (Potapow 2001). An eurytopic species, reaching high abundances in various biotopes (forests, open grasslands). It is also common but less abundant in strongly disturbed areas (intensively cultivated agricultural fields, recultivated sites etc.). Similarly to the Sopron samples, *Cryptopygus bipunctatus* was found among the most dominant species (D = 17%) also in the Fertőrákos sample. Dominant species also include the Palaearctic *Collembola Hypogastrura vernalis* (D = 15,5%), characteristic for seashore meadows with wide distribution along the Nordic coasts. Sporadic European inland records (mostly in continental sands) are also known (Thibaud et al. 2004).

The result of various similarity calculations is presented in *Tables 4-7*. Despite of the high *Collembola* species richness, the similarity between the Sopron-Tómalom-Fertőrákos samples is very low.

Table 4. Renkonen's similarity

	Sopron	Tómalom
Tómalom	6.4028	
Fertőrákos	36.6434	5.1819

Table 5. Sørensen's similarity

	Sopron	Tómalom
Tómalom	0.1276	
Fertőrákos	0.3509	0.1250

Table 6. Jaccard's similarity

	Sopron	Tómalom
Tómalom	6.8182	
Fertőrákos	21.2767	7.1429

Table 7. Wainstein's similarity

	Sopron	Tómalom
Tómalom	43.6557	
Fertőrákos	779.6476	37.0137

The Renkonen similarity index indicates a relatively high similarity between Sopron and Fertőrákos while just the opposite can be observed for the Sopron and Tómalom samples. The other similarity coefficients calculated (Sørensen quotient, Jaccard's species identity index, Weinstein similarity index) also show the "marginal" position of Tómalom.

Table 8 shows the most important structural properties of the *Collembola* communities.

Table 8. Structural properties of *Collembola* communities

	Sopron	Tómalom	Fertőrákos
Species richness (<i>S</i>)	28	20	29
Number of individuals	1245	153	1831
Shannon diversity (<i>H</i>)	1.883	2.628	2.01
Equitability (<i>J</i>)	0.5651	0.8771	0.597

Despite of the lower abundance in Tómalom the species diversity is relatively high, higher than for the other two habitats. The higher community equitability value for the Tómalom sample can be explained with the lower species richness with a relevant implication on the algorithm used.

Table 9 shows the results of *t* test based on Shannon diversity indices.

Table 9. Results (*t*-values) of comparison of Shannon diversity indices (* significant difference at $P=0.05$ level)

	Sopron	Tómalom
Tómalom	5.0778*	
Fertőrákos	1.9089	4.0871*

Concerning the Sopron and Fertőrákos samples, there are no remarkable differences between diversity values of *Collembola* communities collected in these habitats. Significant difference can be observed between the Sopron - Tómalom and the Tómalom - Fertőrákos samples.

CONCLUSION

Knowing the ecology of the collected species a strong connection between them and the moss species cannot be proved. Their presence is supposedly determined by the surrounding habitats.

Typical bryobiont species – like e.g. *Hymenaphorura creatricis* (Traser – Salló 2003) – were not recorded, only some bryophile species (e.g. *Xenylla boernerii*) were found. Among the most dominant *Collembolas* there are no species living solely on moss. Most of the dominant species collected are eurytopic and less sensitive to anthropogenic disturbance.

REFERENCES

- BABENKO, A. B. – CHERNOVA, N. M. – POTAPOV, M. B. – STEBAEVA, M. B. (1994): Collembola of Russia and adjacent countries: Family *Hypogastruridae*. Nauka, Moscow. 336 p. (in Russian)
- BALOGH, J. (1958): Lebensgemeinschaften der Landtiere. Akadémia kiadó, Budapest-Berlin. 560 p.
- BRETFELD, G. (1999): *Symphyleona*. In: DUNGER, W. ed.: Synopses on Palaearctic *Collembola*. Vol. 2. Staatliches Museum für Naturkunde, Görlitz. 318 p.
- DEHARVENG, L. (1982): Cle de détermination des genres de Neanurinae (Collembola) d'Europe et la région Méditerranéenne, avec description de deux nouveaux genres. Université P. Sabatier. Travaux du Laboratoire d'Écobiologie des Arthropodes Edaphiques, Toulouse Vol. 3. Fasc. 4: 7-13.
- DUNGER, W. (1983): Tiere im Boden. A. Ziemsen Verlag, Lutherstadt Wittenberg. 280 p.
- ERZBERGER, P. – PAPP, B. (2004): Annotated checklist of Hungarian bryophytes. *Studia Bot. Hung.* 35: 91 – 149.
- FJELLBERG, A. (1980): Identification keys to Norwegian *Collembola*. Norsk Entomologisk Forening. 152 p.
- FJELLBERG, A. (1998): The Collembola of Fennoscandia and Denmark. Part I.: *Poduromorpha*. *Fauna Entomologica Scandinavica*, Vol.: 35. Brill, Leiden – Boston – Köln. 184 p.
- GISIN, H. (1960): Collembolenfauna Europas. *Museum d'Histoire Naturelle*, Genève. 312 p.
- HEDENAS, L. (1993): Field and microscope keys to the Fennoscandian species of the *Calliergon-Scorpidium-Drepanocladus* complex. *Biodetektor AB*, Marsta, Sweden. 79 p.
- JORDANA, R. – ARBEA, J. I. – CARLOS SIMÓN, M. J. L. (1997): *Collembola Poduromorpha*. *Fauna Iberica*, Vol.: 8. Museo Nacional de Ciencias Naturales, Madrid. 807 p.
- MASSOUD, Z. (1967): Monographie des Neanuridae, Collemboles Poduromorphes a pièces buccales modifiées. Centre National de la Recherche Scientifique, Paris. 399 p.
- ORBÁN S. – VAJDA L. (1983): Magyarország mohafldrájának kézikönyve. Akadémiai Kiadó, Budapest. 518 p.
- PALACIOS-VARGAS, J.G. – CASTANO MENESES, G. – GOMEZ-ANAYA, J.A. (1998): *Collembola* from the canopy of a Mexican Tropical deciduous forest. *Pan-Pacific Entomologist* 74(1): 47-54.
- PALACIOS-VARGAS, J.G. – CASTANO MENESES, G. – PESCADOR RUBIO, A. (1999): Phenology of canopy arthropods of a tropical deciduous forest in Western Mexico. *Pan-Pacific Entomologist* 75(4): 200-211.
- POMORSKI, J. R. (1998): *Onychiurinae* of Poland (*Collembola: Onychiuridae*). Wrocław. 201 p.
- POTAPOV, M. (2001): *Isotomidae*. In: DUNGER, W. ed.: Synopses on Palaearctic *Collembola*. Vol.: 3. *Abhandlungen und Berichte des Naturkundemuseums Görlitz*. Bd. 73/Nr. 2. 603 p.
- SMITH, A. J. E. (1978): The moss flora of Britain and Ireland. Cambridge University Press, Cambridge. 706 p.
- SMITH, A. J. E. (1990): The liverworts of Britain and Ireland. Cambridge University Press, Cambridge. 362 p.
- STACH, J. (1960): The Apterygotan fauna of Poland in relation to the world fauna of this group of insects. Tribe: *Orchesellini*. Państwowe Wydawnictwo Naukowe, Krakowie. 151 p.
- STACH, J. (1963): The Apterygotan fauna of Poland in relation to the world fauna of this group of insects. Tribe: *Entomobryini*. Państwowe Wydawnictwo Naukowe, Krakowie. 126 p.
- THIBAUD, J. M. – SHULZ, H. J. – DA GAMA ASSALINO, M. M. (2004): *Hypogastruridae*. In: DUNGER, W. ed.: Synopses on Palaearctic *Collembola*. Vol.: 4. *Abhandlungen und Berichte des Naturkundemuseums Görlitz*. Bd. 75/Nr. 2. 287 p.
- TRASER, GY. (2002): The *Collembola* of the Fertő-Hanság National Park. In: Mahunka, S. (ed.): The fauna of the Fertő-Hanság National Park. *Magyar Természettudományi Múzeum*. 259-270.
- TRASER, GY. – CHRISTIAN, E. (1992): *Lepidocyrtus peisonis* sp. n., ein neuer Springschwanz aus dem Neusiedlersee-Gebiet (*Collembola: Entomobryidae*). *Folia Entomol. Hung.* 52: 119-121.
- TRASER, GY. – SALLÓ, M. (2004): Új ugróvillás (*Insecta: Collembola*) adatok a Kőszegi-hegységből. [New springtail (*Insecta: Collembola*) data from the Kőszeg Mountains] *Természetvédelmi Közlemények* 11: 229-232. (in Hungarian)
- VARGA, J. – OLDAL, V. (1998): A légszennyezés hatása a mohák állatközösségére. [Effects of air pollution on bryophyl animal communities] *Acta Academiae Paedagogicae Agriensis, Nova Series*, XXII: 83-92. (in Hungarian)

- VARGA, J. – VARGHA, B. (1992): Adatok az Upponyi-szoros Bryofaunájának összetételére vonatkozóan. [Data on the bryofauna of the Uppony Ravine] Acta Academiae Pedagogicae Agriensis, Nova Series, XXI: 83-92. (in Hungarian)
- VARGA, J. (1989) Tájékoztató vizsgálatok különböző stratégiájú mohafajok faunáján. [Primary studies on the fauna of moss species using different ecological strategies] Acta Academiae Paedagogicae Agriensis, Nova Series XIX/IX: 59-72. (in Hungarian)
- VARGA, J. (1991): Adatok Felsőtárkány területének bryofaunájára vonatkozóan. [Data on the bryofauna of Felsőtárkány] Acta Academiae Pedagogicae Agriensis Nova Series XX: 81-88. (in Hungarian)
- VARGA, J. (1992): Analysis of the fauna of protected moss species. Biological Conservation 59: 171-173.
- WEINER, W. M. (1996): Generic revision of *Onychiurinae* (Collembola: Onychiuridae) with a cladistic analysis. Ann. Soc. Entomol. Fr. (N. S.). 32(2): 163-200.
- ZIMDARS, B. – DUNGER, W. (1994): *Tullbergiinae*. In: DUNGER, W. ed.: Synopses on Palaearctic Collembola. Vol.: I. Abhandlungen und Berichte des Naturkundemuseums Görlitz. Bd. 68/Nr. 3-4. 71 p.

Appendix. Raw data of the Collembola specimens collected

Tortella tortuosa (HEDW.) LIMPR.

Fertőrákos, 06. 05. 2004: *Hypogastrura vernalis* 35; *Xenylla brevicauda* 50; *Lepidocyrtus lanuginosus* 1; *Pseudosinella octopunctata* 12; *Orchesella cincta* 2; *Orchesella xerothermica* 2; *Folsomia manolachei* 5; *Cryptopygus bipunctatus* 5; *Entomobrya multifasciata* 4; *Parisotoma notabilis* 11; *Protaphorura armata* 2; *Isotomiella minor* 1;

Pleurochaete squarrosa (BRID.) LINDB.

Fertőrákos, 06. 05. 2004: *Lepidocyrtus lanuginosus* 4; *Folsomia manolachei* 237; *Entomobrya handschini* 7; *Entomobrya nivalis* 8; *Fasciosminthurus strigatus* 1; *Parisotoma notabilis* 9; *Cryptopygus bipunctatus* 5; *Isotoma viridis* 2; *Pseudosinella octopunctata* 13; *Hypogastrura vernalis* 24; *Xenylla brevicauda* 70; *Orchesella cincta* 2; *Megalothorax minimus* 1; *Mesaphorura krausbaueri* 1;

Plagiomnium elatum (BRUCH et SCHIMP.) T. J. KOP.

Tómalom, 10. 09. 2003: *Lepidocyrtus cyaneus* 1; *Lepidocyrtus peisonis* 2; *Friesea truncata* 1; *Supraphorura furcifera* 1;

Plagiomnium undulatum (HEDW.) T. J. KOP.

Sopron, 27. 03. 2003: *Lepidocyrtus lanuginosus* 8; *Sminthurinus elegans* 20; *Mesaphorura macrochaeta* 1; *Parisotoma notabilis* 5; *Heteromurus major* 5; - **Sopron, 10. 09. 2003:** *Parisotoma notabilis* 2; *Lepidocyrtus lanuginosus* 10;

Climacium dendroides (HEDW.) F. WEBER et D. MOHR

Sopron, 10. 09. 2003: *Megalothorax minimus* 4; *Parisotoma notabilis* 4; *Folsomia penicula* 12; *Heteromurus nitidus* 2; *Lepidocyrtus lanuginosus* 5; *Cryptopygus bipunctatus* 42; *Pseudosinella alba* 2;

Thuidium philibertii LIMPR.

Sopron, 20. 03. 2003: *Heteromurus major* 1; *Tomocerus minor* 1; *Metaphorura affinis* 1; *Parisotoma notabilis* 35; *Orchesella cincta* 2; *Cryptopygus bipunctatus* 20; *Sminthurinus elegans* 6; *Megalothorax minimus* 2; *Pseudosinella alba* 2;

Thuidium abietinum (HEDW.) SCHIMP.

Fertőrákos, 06. 05. 2004: *Mesaphorura krausbaueri* 2; *Lepidocyrtus lanuginosus* 9; *Cryptopygus bipunctatus* 1; *Hypogastrura vernalis* 48; *Folsomia manolachei* 41; *Megalothorax minimus* 1; *Entomobrya nigriventris* 3; *Fasciosminthurus strigatus* 3; *Pseudosinella octopunctata* 2; *Isotoma viridis* 2; *Parisotoma notabilis* 10; *Heteromurus major* 1; *Willemia virae* 1; *Heterosminthurus bilineatus* 2; *Entomobrya handschini* 4; *Isotomiella minor* 1;

Campylium chrysophyllum (BRID.) LANGE

Fertőrákos, 06. 05. 2004: *Cryptopygus bipunctatus* 1; *Lepidocyrtus lanuginosus* 2; *Parisotoma notabilis* 8; *Hypogastrura vernalis* 4;

Amblystegium varium (HEDW.) LINDB.

Tómalom, 10. 09. 2003: *Lepidocyrtus cyaneus* 1; *Lepidocyrtus peisonis* 5; *Friesea truncata* 3; *Megalothorax minimus* 5; *Spaeridia pumilis* 5; *Isotomiella minor* 2; *Supraphorura furcifera* 2; *Parisotoma notabilis* 2; *Lepidocyrtus paradoxus* 1; *Neanura muscorum* 1;

Drepanocladus cossonii (SCHIMP.) LOESKE

Tómalom, 10. 09. 2003: *Lepidocyrtus peisonis* 3; *Lepidocyrtus paradoxus* 1; *Brachystomella parvula* 2; *Isotomurus prasinus* 8; *Supraphorura furcifera* 1; *Friesea truncata* 2; *Megalothorax minimus* 2; *Spaeridia pumilis* 1; *Isotomiella minor* 1;

Calliergonella cuspidata (HEDW.) LOESKE

Tómalom, 10. 09. 2003: *Lepidocyrtus cyaneus* 1; *Lepidocyrtus peisonis* 1; *Isotomurus prasinus* 5; *Friesea truncata* 4; *Megalothorax minimus* 6; *Spaeridia pumilis* 18; *Isotomiella minor* 10; *Supraphorura furcifera* 3; *Parisotoma notabilis* 5; *Pachyotoma crassicauda* 2;

Brachystomella parvula 1; *Neelides minutus* 2; *Micranurida pygmaea* 3; *F. quadrioculata* 4; *Dicyrtoma fusca* 2; - **Sopron, 06. 03. 2003:** *Heteromurus major* 3; *Lepidocyrtus lanuginosus* 2; *Parisotoma notabilis* 5; *Cryptopygus bipunctatus* 220; *Pseudosinella alba* 1; - **Sopron, 20. 03. 2003:** *Lepidocyrtus lanuginosus* 20; *Heteromurus major* 4; *Tomocerus cf. baudoti* 1; *Cryptopygus bipunctatus* 100; *Parisotoma notabilis* 200; *Spaeridia pumilis* 4; *Mesaphorura hylophila* 12; *Doutnacia xerophila* 6; *Megalothorax minimus* 2; *Orchesella cincta* 2; *Heteromurus major* 8; *Sminthurinus elegans* 2; - **Sopron, 10. 09. 2003:** *Megalothorax minimus* 1; *Spaeridia pumilis* 2; *Parisotoma notabilis* 5; *Folsomia penicula* 6; *Sminthurinus aureus* 2; *Tomocerus minor* 1; *Heteromurus nitidus* 1; *Lepidocyrtus lanuginosus* 6; *Cryptopygus bipunctatus* 2;

Homalothecium lutescens (HEDW.) H. ROB

Fertőrákos, 06. 05. 2004: *Mesaphorura critica* 4; *Folsomia manolachei* 73; *Parisotoma notabilis* 14; *Heteromurus major* 2; *Hypogastrura vernalis* 64; *Fasciosminthurus strigatus* 2; *Oncopodura crassicornis* 1; *Orchesella cincta* 8; *Orchesella xerothermica* 1; *Cryptopygus bipunctatus* 11; *Mesaphorura krausbaueri* 3; *Parisotoma notabilis* 4; *Lepidocyrtus lignorum* 2; *Pseudosinella octopunctata* 1;

Brachythecium rutabulum (HEDW.) SCHIMP.

Tómalom, 10. 09. 2003: *Lepidocyrtus peisonis* 4; *Isotomurus prasinus* 8; *Spaeridia pumilis* 1; *Pachytoma crassicauda* 4;

Scleropodium purum (HEDW.) LIMPR.

Sopron, 10. 09. 2003: *Parisotoma notabilis* 8; *Folsomia penicula* 11; *Lepidocyrtus lanuginosus* 10;

Eurhynchium hians (HEDW.) SANDLE LAC.

together with *Scleropodium purum* (HEDW.) LIMPR. and with *Brachythecium rutabulum* (HEDW.) SCHIMP. samples **Sopron, 06. 03. 2003:** *Mesaphorura macrochaeta* 2; *Lepidocyrtus lanuginosus* 17; *Heteromurus major* 4; *Orchesella cincta* 6; *Isotomurus cf. palustris* 6; *Parisotoma notabilis* 35; *Cryptopygus bipunctatus* 47; *Megalothorax minimus* 2; *Pseudosinella alba* 1;

Hypnum cupressiforme HEDW.

Fertőrákos, 06. 05. 2004: *Pseudosinella alba* 1; *Pseudosinella octopunctata* 3; *Lepidocyrtus lanuginosus* 7; *Entomobrya nivalis* 3; *Folsomia manolachei* 270; *Parisotoma notabilis* 93; *Hypogastrura vernalis* 32; *Megalothorax minimus* 2; *Mesaphorura krausbaueri* 23; *Cryptopygus bipunctatus* 58; *Xenylla boernerii* 145; *Cryptopygus bipunctatus* 88; *Heterosminthurus bilineatus* 2; - **Sopron, 02. 03. 2004:** *Xenylla boernerii* 400; *Hypogastrura socialis* 4; *Entomobrya nivalis* 1; *Lepidocyrtus violaceus* 1; *Entomobrya corticalis* 2; *Orchesella bifasciata* 1; *Xenylla brevicauda* 2; *Xenylla maritima* 1; *Cyphoderus albinus* 1;

Rhytidium rugosum (HEDW.) KINDB.

Fertőrákos, 06. 05. 2004: *Cryptopygus bipunctatus* 150; *Hypogastrura vernalis* 75; *Lepidocyrtus lanuginosus* 2; *Mesaphorura krausbaueri* 8; *Parisotoma notabilis* 10; *Cyphoderus albinus* 1; *Folsomia manolachei* 6; *Fasciosminthurus strigatus* 1;

Rhytidiadelphus triquetrus (HEDW.) WARNST.

Sopron, 20. 03. 2003: *Tomocerus minor* 1; *Tomocerus cf. baudoti* 2; *Lepidocyrtus lanuginosus* 8; *Folsomia penicula* 2; *Heteromurus major* 5; *Parisotoma notabilis* 10; *Sminthurinus elegans* 5