

New Reports of Dothistroma Needle Blight in Eurasian Countries

Irene BARNES^{a*} – Thomas KIRISITS^b – Alexander AKULOV^c – D. B. CHHETRI^d –
Michael J. WINGFIELD^a – Timur BULGAKOV^e – Brenda D. WINGFIELD^a

^aForestry and Agricultural Biotechnology Institute, Department of Genetics, University of Pretoria, *South Africa*;

^bInstitute of Forest Entomology, Forest Pathology and Forest Protection, Department of Forest and Soil Sciences, University of Natural Resources and Applied Life Sciences, Vienna, Austria.

^cDepartment of Mycology and Plant Resistance, V.N. Karasin National University of Kharkiv, Kharkiv, Ukraine

^dRenewable Natural Resources Research Centre, Jakar, East Central Region, Council for RNR Research of Bhutan, Ministry of Agriculture, Royal Government of Bhutan, Bhutan

^eDepartment of Biology and Pedology, South Federal University, Rostov-na-Donu, Russia.

Extended abstract – One of the most serious needle diseases that affect pines (*Pinus* spp.) is Dothistroma needle blight (DNB). Two species of fungi are responsible for causing this disease (Barnes et al. 2004). These are *Dothistroma septosporum* (teleomorph: *Mycosphaerella pini*) that has a worldwide distribution and infects a wide range of *Pinus* spp. and *D. pini* (teleomorph unknown), which has thus far been reported only from the North-Central U.S.A. on the non-native *Pinus nigra* (Barnes et al. 2004). In recent years, there have been increasing numbers of reports of DNB from new hosts and new geographic regions of the Northern Hemisphere (Bradshaw 2004, Bednářová et al. 2006). Moreover, there has been an increase in the intensity of this disease in some parts of Europe and North America (Koltay 2001, Aumonier 2002, Brown et al. 2003, Jankovský et al. 2004, Woods et al. 2005).

Since 2004, we have conducted surveys and inspections of trees in Austria, Bhutan, Hungary, Ukraine and South-Western Russia. These have helped to document DNB on several native and non-native pine species, and to unmask its presence in situations where disease symptoms and signs were not obvious or not typical (Barnes et al. 2007). In 2004, non-native *Pinus peuce* trees in an arboretum in Vienna (Austria) were found to suffer from DNB. In 2005, a non-native *Pinus radiata* tree in Western Bhutan was found with typical DNB symptoms. Further east in Central Bhutan, native *Pinus wallichiana* trees in conifer forests at high elevations, had needle blight symptoms atypical of DNB. These, and other pine needle collections from Hungary, Ukraine and South-Western Russia, with typical DNB symptoms formed the basis of this study.

Isolates from the various hosts and countries were examined morphologically and compared using DNA sequence data. Conidia for all collections were elongated, straight to slightly curved, hyaline and they had one to five septa. Conidial dimensions varied considerably when measured from conidiomata produced on needles and in culture. Detailed measurements indicated that *D. pini* has slightly wider conidia than *D. septosporum*, as has

* Corresponding author: Irene.barnes@fabu.up.ac.za; 74 Lunnon Road, Hillcrest, Pretoria 0002, South Africa.

previously been shown by Barnes et al. (2004). However, these differences are so small that distinguishing *D. septosporum* from *D. pini* based on morphology is virtually impossible.

Comparisons of DNA sequence data for the ITS region of the rDNA and parts of the β -tubulin gene region, were used to unambiguously identify the *Dothistroma* pathogens collected from symptomatic needles. *Pinus peuce* from Austria, *P. radiata* and *P. wallichiana* from Bhutan as well as *Pinus mugo* from Hungary were all found to be infected by *D. septosporum*. In contrast, *Pinus pallasiana* from the Kherson region in Ukraine and the nearby Rostov region in South-Western Russia were infected by *D. pini*. These results represent a new host record for *D. septosporum* on *P. peuce* and a new country report for *D. septosporum* from Bhutan. They also confirm that *D. septosporum* occurs in Hungary. Results also provide a new host report for *D. pini* on *P. pallasiana* and new country reports for *D. pini* in Ukraine and South-Western Russia. The latter records represent the only reports of *D. pini* from outside the North-Central U.S.A. The new host and country records provided here are consistent with the continuing trend of reports of the DNB pathogens from new hosts and new geographic areas during the last two decades, particularly in the Northern Hemisphere.

***Dothistroma septosporum* / *Dothistroma pini* / *Mycosphaerella pini* / Bhutan / Russia / Ukraine**

REFERENCES

- AUMONIER, T. (2002): La maladie des bandes rouges toujours en augmentation en [Dothistroma needle blight (*Dothistroma septospora*) still on the increase in 2001]. Les Cahiers du DSF, 1-2002 (La Santé des Forêts [France] en 2000 et 2001), Min. Agri. Alim. Pêche Aff. rur. (DERF), Paris, 58-60.
- BARNES, I. – CROUS, P.W. – WINGFIELD, M.J. – WINGFIELD, B.D. (2004): Multigene phylogenies reveal that red band needle blight of *Pinus* is caused by two distinct species of *Dothistroma*, *D. septosporum* and *D. pini*. Stud. Mycol. 50: 551-565.
- BARNES, I. – KIRISITS, T. – AKULOV, A. – CHHETRI, D.B. – WINGFIELD, B.D. – BULGAKOV, T.S. – WINGFIELD, M.J. (in press): New host and country records of the *Dothistroma* needle blight pathogens from Europe and Asia. For. Path.
- BEDNÁŘOVÁ, M. – PALOVČÍKOVÁ, D. – JANKOVSKÝ, L. (2006): The host spectrum of *Dothistroma* needle blight *Mycosphaerella pini* E. Rostrup – new hosts of *Dothistroma* needle blight observed in the Czech Republic. J. For. Sci. 52: 30-36.
- BRADSHAW, R.E. (2004): *Dothistroma* (redband) needle blight of pines and the dothistromin toxin: a review. For. Path. 34: 163-185.
- BROWN, A. – ROSE, D. – WEBBER, J. (2003): Red Band Needle Blight of Pines. Forestry Commission Edinburgh: Information Note.
- JANKOVSKÝ, L. – BEDNÁŘOVÁ, M. – PALOVČÍKOVÁ, D. (2004): *Dothistroma* needle blight *Mycosphaerella pini* E. Rostrup, a new quarantine pathogen of pines in the CR. J. For. Sci. 50: 319-326.
- KOLTAY, A. (2001): A *Dothistroma septospora* (Dorog.) Morlet előfordulása a hazai feketefenyő (*Pinus nigra* Arn.) állományokban, és az ellene alkalmazott vegyszeres védekezési kísérletek eredményei [Incidence of *Dothistroma septospora* (Dorog.) Morlet in the Austrian pine (*Pinus nigra* Arn.) stands in Hungary and results of chemical control trials]. Növényvédelem. 37: 231-235.
- WOODS, A. – COATES, K.D. – HAMANN, A. (2005): Is an unprecedented *Dothistroma* needle blight epidemic related to climate change? BioScience 55: 761-769.